

# CHLOROPHYLL *A* CONCENTRATION OF FRESH WATER PHYTOPLANKTON ANALYSED BY ALGORITHMIC BASED SPECTROSCOPY

**FAIROOZ BINTI JOHAN**

**UNIVERSITI SAINS MALAYSIA**

2016

**CHLOROPHYLL *A* CONCENTRATION OF  
FRESH WATER PHYTOPLANKTON ANALYSED  
BY ALGORITHMIC BASED SPECTROSCOPY**

**by**

**FAIROOZ BINTI JOHAN**

**Thesis submitted in fulfillment of the requirements  
for the degree of Doctor of Philosophy**

**Februari 2016**

## ACKNOWLEDGEMENTS

I thanks to Allah s.w.t. for being able to finish my PhD thesis after having almost three years time. I faced a lot of different challenge while i carried out my research work and thesis writing but finally i managed to finish with Allah s.w.t will.

First and foremost, i would like to express my sincere gratitude to my main supervisor, Prof. Dr Mohd Zubir bin Mat Jafri for always been there, supporting and helping me throughout my research. He never hesitated to share ideas and guide me whenever possible, even for answering some of my unintelligent questions about this research. I am much indebted to him for spending time to read this thesis and gave critical comment about it.

Besides, I would like to thank to my co-supervisor, Assoc. Prof. Dr Lim Hwee San for his contribution and guidance especially correcting my mistake in the research and using so much time to read this thesis and gave critical comment about it. Not forgetting my co-supervisor from School of Biological Science, Assoc. Prof Dr Wan Maznah binti Wan Omar for giving me permission to use the plankton lab for my research purpose. Moreover, she was kind enough to teach me with patience about phytoplankton when this topic was once unfamiliar to me.

I would like to include a special note of thanks to scholarship and grants for giving me financial support to finish this qualification. Without this scholarship and grants, i would not be able to continue my journey in PhD.

- i. Department of Higher Education (MyPhD Scholarship).
- ii. Research University grant no: 1001/PFIZIK/811220
- iii. Fundamental Research Grant no: 203/PFIZIK/6711349

To my husband, Mohd Afendi bin Abdul Mutalib, thank you for always being supportive and patient with me in my PhD journey. To my kids, Nur Fadhia Adriana and Fuzail Aryan, who always sent me off to work at the door every morning and waiting for me to come home every evening, thank you for bearing with me. To my family, especially my mother, Masitah binti Sidek, my mother in law, my sibling (Firdaus, Faisal Faliq, Faramalena and Fitri Aildil), my sisters and brothers in law thank you for being so understanding throughout my three years of study. I would like to dedicate this thesis to my late father, Johan bin Abu Hassan. Thanks for being a good father. Although you were gone six years, you are never to be forgotten.

A special thanks to all staffs from School of Physics and School of Biological Science for assisting me in related issues throughout my PhD study in Universiti Sains Malaysia. A very thank you to my friends and my lab mates from School of Physics, for lending a helping hand during my research work. Lastly, thank you very much to my friend from School of Biological accompanying while i took water samples in Tasik Harapan. I really appreciate it.

## **TABLE OF CONTENTS**

	<b>Page</b>
<b>ACKNOWLEDGEMENTS</b>	<b>ii</b>
<b>TABLE OF CONTENTS</b>	<b>iii</b>
<b>LIST OF TABLES</b>	<b>vii</b>
<b>LIST OF FIGURES</b>	<b>viii</b>
<b>LIST OF SYMBOL AND ABBREVIATIONS</b>	<b>x</b>
<b>ABSTRAK</b>	<b>xi</b>
<b>ABSTRACT</b>	<b>xii</b>
<b>CHAPTER 1 - INTRODUCTION</b>	
1.1 Background	1
1.2 Spectroscopy	4
1.3 Problem Statement	5
1.4 Research Objectives	6
1.5 Research Scope	6
1.6 Research Location	6
1.7 Novelty of Study	7
1.8 Structure of the Thesis	7
<b>CHAPTER 2 - LITERATURE REVIEW</b>	
2.1 Introduction	8
2.2 Definition and Background of Phytoplankton	9
2.2.1 Green Algae	14
2.2.2 Cyanobacteria	14
2.2.3 Red algae	14

2.2.4	Golden algae/Diatom	15
2.2.5	Brown algae	15
2.2.6	Coccolithophores	16
2.2.7	Dinoflagellates	16
2.2.8	Yellow-green algae	16
2.2.9	Euglenids	17
2.2.10	Cryptomonads	17
2.3	Chlorophyll	18
2.4	Chlorophyll <i>a</i>	19
2.5	Other Chlorophyll	21
2.6	Methods to Measure Chlorophyll	22
2.7	Phytoplankton Reflectance	24
2.8	Optical Properties of Water	29
2.9	Technological Application of Phytoplankton	33
2.10	Summary	35

## **CHAPTER 3 - DEVELOPMENT OF OPTICAL MODEL FOR PHYTOPLANKTON IN FRESH WATER**

3.1	Introduction	36
3.2	Regression Algorithm	36
3.3	Summary	42

## **CHAPTER 4 - METHODOLOGY**

4.1	Introduction	43
4.2	Study Area	43

4.3	Methodology and Instruments	45
4.3.1	Random Sampling Technique	45
4.3.2	Cultured Method	47
4.3.3	Reflectance Measurement	49
4.3.4	Extraction Chlorophyll <i>a</i> Concentration	49
4.3.5	Chlorophyll <i>a</i> Concentration Measurement	54
4.3.6	Plankton net	55
4.3.7	Global positioning System	56
4.3.8	Spectroradiometer	56
4.3.9	ViewSpec Pro Software	57
4.4	Summary	58

## **CHAPTER 5 - RESULT AND DISCUSSION**

5.1	Introduction	60
5.2	Reflectance Properties of Phytoplankton in Freshwater	61
5.3	Correlation between Absorbance and Chlorophyll <i>a</i> Concentration	69
5.4	Correlation between reflectance and chlorophyll <i>a</i> concentration	72
5.5	Validation The Concentration Of Chlorophyll <i>a</i> from The Algorithms Developed.	77
5.5.1	Validation for Single Wavelength	77
5.5.2	Validation for Two Wavelengths	78
5.5.3	Validation for Three Wavelengths	79
5.5.4	Discussion	80
5.5.5	Summary	81

<b>CHAPTER 6 - CONCLUSION AND FUTURE RECOMMENDATIONS</b>	
6.1 Conclusion	83
6.2 Future Recommendation	85
<b>REFERENCE</b>	86
<b>APPENDIX A</b> The volumes, surface area and Surface Area / Volume ratio of phytoplankton species	97
<b>APPENDIX B</b> Vegetation spectral reflectance characteristics in specific regions of electromagnet spectrum	99
<b>APPENDIX C</b> Predominant photosynthetic pigment, storage product, and cell wall component of the major algal group	103
<b>APPENDIX D</b> FieldSpec® HandHeld 2 Specifications	104
<b>APPENDIX E</b> Regression Analysis and Graphs from software	105
<b>LIST OF PUBLICATION</b>	111

## LIST OF TABLES

	<b>Page</b>
<b>Table 2.1:</b> Type of plankton base on habitat	10
<b>Table 5.1:</b> Wavelength range for VIS and NIR ranges	65



## LIST OF FIGURES

	<b>Page</b>
<b>Figure 1.1 :</b> Size and types of phytoplankton	3
<b>Figure 1.2 :</b> Spectrum of electromagnet	5
<b>Figure 2.1 :</b> Chart of Phytoplankton Group	13
<b>Figure 2.2 :</b> Scatterplots of reflectance ratio versus chlorophyll <i>a</i>	23
<b>Figure 2.3:</b> Reflectance spectra of water and phytoplankton	27
<b>Figure 2.4:</b> Typical reflectance sensitivities as controlled by pigment, cell structure and water content	27
<b>Figure 4.1:</b> Study area	44
<b>Figure 4.2:</b> Tasik Harapan	45
<b>Figure 4.3:</b> Refrigerator that used to store the samples	46
<b>Figure 4.4:</b> Autoclave	47
<b>Figure 4.5:</b> Preparing sample place	48
<b>Figure 4.6:</b> Culture room	48
<b>Figure 4.7:</b> Setting the instrument	49
<b>Figure 4.8:</b> Filtering of water sample using vacuum pump	50
<b>Figure 4.9:</b> Step to fold the filter paper and keep in refrigerator	51
<b>Figure 4.10:</b> Blended step, using Heidolph shaker type Reax 2000	52
<b>Figure 4.11:</b> Blended step, using manual step	52
<b>Figure 4.12:</b> Centrifuge tubes containing chlorophyll	53
<b>Figure 4.13:</b> A centrifuge	53
<b>Figure 4.14:</b> Preparing to bring to Biochemistry lab	54
<b>Figure 4.15:</b> Spectrophotometer	55
<b>Figure 4.16:</b> Phytoplankton net	56
<b>Figure 4.17:</b> Spectroradiometer	57

<b>Figure 4.18:</b>	Example showing a spectral graph ASD	58
<b>Figure 5.1:</b>	Graphs of reflectance versus wavelength	65
<b>Figure 5.2:</b>	Highest and lowest graphs of reflectance versus wavelength	66
<b>Figure 5.3 :</b>	1st derivative reflectance of phytoplankton	67
<b>Figure 5.4:</b>	Absorbance and concentration of chlorophyll <i>a</i> for 630 nm	70
<b>Figure 5.5:</b>	Absorbance and concentration of chlorophyll <i>a</i> for 647 nm	70
<b>Figure 5.6:</b>	Absorbance and concentration of chlorophyll <i>a</i> for 664 nm	71
<b>Figure 5.7:</b>	Concentration of chlorophyll <i>a</i> versus reflectance of phytoplankton for 438 nm	73
<b>Figure 5.8:</b>	Concentration of chlorophyll <i>a</i> versus reflectance of phytoplankton for 550 nm	74
<b>Figure 5.9:</b>	Concentration of chlorophyll <i>a</i> versus reflectance of phytoplankton for 675 nm	75
<b>Figure 5.10:</b>	Relationship between predicted and actual concentration of chlorophyll <i>a</i> for single wavelength	78
<b>Figure 5.11:</b>	Relationship between predicted and actual concentration of chlorophyll <i>a</i> for two wavelengths	79
<b>Figure 5.12:</b>	Relationship between predicted and actual concentration of chlorophyll <i>a</i> for three wavelengths	80

## LIST OF SYMBOLS AND ABBREVIATIONS

BBM	Bold's Basal Medium
Chl <i>a</i>	Chlorophyll <i>a</i>
NIR	Near infrared
VIS	Visible
<b>R</b>	Reflectance spectrum
R <sub>s</sub>	Reflectance of water surface
R <sub>rs</sub>	Reflectance for remote sensing
R <sub>L</sub>	Light reflectance
<i>R</i>	Coefficient
C <sub>a</sub>	Concentration of chlorophyll <i>a</i>
µg/L	microgram per litres
mg/L	milligram per litres
λ	Wavelength

# **KEPEKATAN KLOOROFIL *A* FITOPLANKTON AIR TAWAR DIANALISIS DENGAN SPEKTROSKOPI BERASASKAN ALGORITMA**

## **ABSTRAK**

Fitoplankton ialah tumbuhan mikroskopik sel tunggal yang memainkan peranan penting di dalam ekosistem sebagai pengeluar primari utama melalui aktiviti fotosintesis. Objektif utama kajian ini ialah untuk mengkaji pencirian melalui hubungan pantulan dan kepekatan klorofil *a* fitoplankton di dalam air tawar. Melalui, kajian ini, pantulan fitoplankton diambil dengan menggunakan spektrometri untuk melihat hubungan antara pantulan dan panjang gelombang fitoplankton di dalam air tawar. Satu lokasi tasik dalam Universiti Sains Malaysia yang dikenali sebagai Tasik Harapan dipilih untuk kajian ini. Sebanyak 20 liter sampel air diambil dan ditapis dengan menggunakan jaring fitoplankton. Sampel air yang diambil dianalisis di dalam makmal untuk menentukan pantulan dan kandungan kepekatan klorofil-*a* fitoplankton. Dua jenis spektrometer digunakan di dalam kajian ini pertama, spektrometri digunakan untuk mengukur pantulan fitoplankton dan kedua, spektrofotometer digunakan untuk mengukur kepekatan klorofil *a*. Sampel air yang diambil dikultur terlebih dahulu dengan memasukkan dalam medium yang dikenali sebagai Bold's Basal sebelum nilai pantulan diambil agar pengukuran pantulan tepat ke atas fitoplankton. Algoritma-algoritma yang menghubungkan kepekatan klorofil *a* dengan pantulan fitoplankton digunakan. Keputusan yang baik diperolehi di dalam kajian ini yang dibuktikan oleh nilai  $R^2$  yang baik di dalam

analisis keputusan melalui algoritma yang dibangun. Tiga kawasan jalur yang difokuskan di dalam kajian ini ialah jalur merah, jalur hijau dan jalur biru. Jalur-jalur ini sesuai digunakan untuk menganalisis fitoplankton. Nilai pantulan bagi setiap jalur ditentukan merujuk kepada kepekatan klorofil *a* untuk kalibrasi algoritma. Selain itu, pelbagai nilai panjang gelombang diuji dan nilai  $R^2$  dibandingkan. Akhirnya, tiga panjang gelombang dipilih iaitu 438 nm, 550 nm dan 675 nm. Pemilihan tiga jarak gelombang ini adalah kerana gelombang-gelombang sepadan dengan fitoplankton dan klorofil *a*. Lagipun nilai  $R^2$  yang diperolehi juga sesuai dengan menggunakan algoritma yang dibangun di dalam kajian ini. Pengesahan penggunaan algoritma juga dilakukan di tasik yang sama tetapi tarikh pengambilan sampel air tidak sama. Keputusan pengesahan ini memberi keputusan yang sangat baik dengan nilai  $R^2$  yang tinggi. Hasil kajian daripada keputusan tasik ini, menunjukkan algoritma yang dibangun untuk mengukur kepekatan klorofil *a* fitoplankton berjaya. Selain itu, beberapa cadangan juga diberikan untuk memperbaiki kajian ini pada masa akan datang.

# **CHLOROPHYLL *a* CONCENTRATION OF FRESH WATER PHYTOPLANKTON ANALYSED BY ALGORITHMIC BASED SPECTROSCOPY**

## **ABSTRACT**

Phytoplankton are microscopic single-celled plants that play an important role in the ecosystem as a major primary producers through photosynthesis. The main objective of this study was to investigate properties through reflection and chlorophyll *a* concentrations of phytoplankton in freshwater. In this study, reflection phytoplankton were taken using spectroradiometer to observe the relationship between the reflectance and the wavelengths of phytoplankton in freshwater. The lake selected for this study is known as Tasik Harapan located in Universiti Sains Malaysia. In this study, 20 litres water samples undergone the filtering using phytoplankton net. The water samples collected were analyzed in the laboratory to determine the reflectance and chlorophyll *a* concentration of phytoplankton. Two spectrometers were used in this study, firstly, spectroradiometer which was used to measure the reflectance of phytoplankton and secondly, spectrophotometer which was used to measure the concentration of chlorophyll *a*. The water samples taken were prior to culture in the medium known as Bold's Basal Medium before the reflections were taken so that accurate reflection measurements on phytoplankton can be acquired. The algorithms that related between chlorophyll *a* concentration and reflectance of phytoplankton were used. The good results are obtained in this study

which are evidenced by the good correlation in the analysis results using the developed algorithm. Three regions were focused which were the red, green and blue. These bands have been identified to be appropriated in analyzing phytoplankton. Reflectance of each band specified referred to the concentration of chlorophyll *a* for calibration algorithm. The wavelength range was tested and the  $R^2$  were compared. Finally, three wavelengths of 438 nm, 550 nm and 675 nm were selected. The selection of these three wavelengths were found to be strongly correlated to phytoplankton and chlorophyll *a*. The value of  $R^2$  obtained also in accordance with the developed algorithm in this study. The water sampling for validation was also taken from the same lake but not on the same date and was analysed using the same algorithm. The validation results showed very good results with a high  $R^2$  value was obtained. The results from this lake proved that the algorithm developed was successful in measuring the concentration of chlorophyll *a* in phytoplankton. In addition, some suggestions are given to repair this study in the future.

## **CHAPTER 1**

### **INTRODUCTION**

#### **1.1 Background**

Fresh water is natural occurring water on the surface of the Earth, like lake, pond and river. Fresh water ecosystem is rich with variety of organism including phytoplankton species. Tasik Harapan is one of the fresh water ecosystem that can be found in Universiti Sains Malaysia (USM) Penang, Malaysia. Fresh water has characteristics of low concentrations of soluble salts and other dissolved solids. Fresh water term does not have the same meaning with potable water. Many surface fresh water and ground water are not suitable for drinking. Besides, fresh water is a natural resource that is vital to the ecosystem life whereas good water quality is required for many ecosystems service (Meybeck & Helmer 1996). From Carpenter et al. (2011), in most parts of the world have suffered fresh water ecosystems degradation seriously. This degradation is strongly related to surrounding land use types reflecting undergoing activities (Perry & Vanderklein, 1996). Consequently, the study on aquatic communities (phytoplankton) and water quality become popular since the last decades (Stomp et al., 2011). Understanding functionality of relationship between ecological lake and watershed changes is a very important step for the effective, long-term conservation or management strategies in selection and application (Silva et al., 2011). Hence, the scientists have used biological indicator for detecting potential changes of biotic communities structure caused by spatial and temporal scale ( Burns & Galbraith, 2007; Van Egeren et al., 2011).




About 70% of the Earth is constituted by water, in which phytoplankton is prevailing organism which is representing the set of autotrophic photosynthetic organisms present in plankton. Phytoplankton is one of the aquatic organisms in fresh water ecosystem and primary biotic community indicating changes in ecological (Padisák et al., 2006). Phytoplankton is also a microscopic plant that is basis of food chain for organisms in the ecosystem (Nyananyo et al 2006). Besides, it is also important as implication in the regulation of marine ecosystems, for examples ecotones between terrestrial, fresh water, and marine habitats. Other functions of phytoplankton concern the global climatic changes control by the fixing of carbon dioxide excess and regulating the biogenic emissions of sulfur on worldwide scale. Phytoplankton need to stay on the water surface to absorb sunlight for photosynthesis activity. Phytoplankton are influenced by several physical parameter such as temperature, nutrient, water current, pH, conductivity, dissolve oxygen and others (Shamsudin, 1991). These physical parameters are important in phytoplankton distribution determining. In addition, the distribution of phytoplankton are also dependent on spatial and seasonal characteristic in the area (Wong and Wong, 2004). Phytoplankton are also beneficial as ecological indicator for assessment of water quality and ecosystem health (Sagert et al 2008, Webber et al 2005).

There are several reasons why phytoplankton community can be used in assessing water quality as described below (Reynolds et al., 1993; Whitaker et al., 2003; Ekwu et al., 2006; Soininen et al., 2007):

- a) The incident of specific of phytoplankton could be limited primary by abiotic conditions such as lake depth and biotic recent. Abiotic and biotic factors that influenced the abundant and distribution of phytoplankton in fresh water.

- b) The biogeographical studies have showed an increase in the issue of distance decay patterns in microorganism community composition in recent evidence.

Phytoplankton species are proposed to reflect trophic conditions (Rajo et al., 2000). Trophic is means nutrition or growth. For example, in eutrophic fresh waters, the relative importance of chrysophytes to biomass or biodiversity (community structure) decrease but cyanobacteria increase (Watson et al., 1997). The phytoplankton size is the most important single characteristic affecting the ecology. When the phytoplankton become bigger, their volume increase their radii while their surface area arise in proportion to only the square of radius. Figure 1.1 shows the size and types of phytoplankton, meanwhile in **Appendix A** shows the volumes, surface area and surface area per volume ratio of phytoplankton species ( Reynolds 2006).



**Figure 1.1** : Size and types of phytoplankton, (G.M, 1950, Freshwater Algae of the United State)


## 1.2 Spectroscopy

Spectroscopy is used in studying of absorption, emission and other radiation (ie. scattering) by matter which is related to the processes on the wavelength of the radiation. In addition, spectroscopic techniques are widely used in almost all technical field of science and technology including in phytoplankton study. However, spectroscopic techniques are very sensitive and must be set up carefully in study to get an accurate result (Stoner, 2014).

Spectroscopy has a various type and each type shows pictures and characteristics spectrum of matter differently. The variation in intensity of the radiation as a function of the wavelength or frequency showing in a graph is called spectrum (Fusino, 2009). The different regions in the electromagnetic spectrum such as ultraviolet (UV), visible (VIS) and infrared (NIR) are dependent on the matter characteristics. In the UV region, the wavelength ranges from 200 nm to 400 nm and in the VIS region, the ranges of wavelength between 380 nm until 750 nm whereas in the NIR region, the wavelength ranges from 750 nm to 2500 nm. Figure 1.2 shows the electromagnetic spectrum ranges from radiowaves to gamma ray.

The benefits of spectroscopy technique that used in this study are fast and easy. The different ranges of wavelength are very important to know because the comprehensive information can be found by these specific ranges for instant, the characteristics of the biochemical composition of a sample. Usually, the spectroscopy analysis is used for reflectance, absorption and transmission measurement of matter. These instruments measure the amount of energy reflected from the ground or object over different wavelengths (Milton, 1987). These

measurement can be converted to spectral radiance value if suitable equipment calibration factors are valid. The magnitude of spectral is depended on radiance solar incoming amount.



**Figure 1.2** : Spectrum of electromagnet (klug & Cummings, 1997)

### 1.3 Problem Statement

Measuring chlorophyll *a* concentration is a step in the process of monitoring for nuisance phytoplankton blooms that may influence the taste and odor of drinking water sources. These blooms may actually create conditions that are toxic to human, fish, wildlife and livestock. Bodies of water used as drinking water source are also monitored for phytoplankton concentration for the early detection of phytoplankton blooms to minimize filtration system clogs. However, measuring chlorophyll *a* concentration is not simple and time consuming due to the need of sample preparation and may require complicated preparation.

#### **1.4 Research Objectives**

The objectives of this study are

1. To investigate the optical properties of phytoplankton.
2. To find the correlation between reflectance and chlorophyll *a* concentration of phytoplankton.
3. To develop a new algorithm for calculating chlorophyll *a* concentration of phytoplankton in fresh water.
4. To estimate the concentration of phytoplankton from the algorithm developed.

#### **1.5 Research Scope**

This research is mainly focus on optical sensor algorithms to measure concentration chlorophyll *a* of phytoplankton in fresh water. Images and data of phytoplankton will be analysed using the theoretical and statistical methods. Some software also used in analysis.

#### **1.6 Research Location**

In this research, 40 water samples were taken at Tasik Harapan, Universiti Sains Malaysia (USM) Penang around 10 o'clock morning. These water samples were collected to measure the chlorophyll *a* concentration of phytoplankton in this lake area by using spectroradiometer method. The coordinates of study area is 05° 35' N, 100° 29' E with width approximately about 8000 m<sup>2</sup>. This location was chosen because it was very near and therefore easier to make observation frequently.

## **1.7 Novelty of Study**

Studying the spectrum of phytoplankton in fresh water provides the new algorithms of measuring concentration of chlorophyll *a* in suitable wavelengths for phytoplankton. The concentration of the sample can be calculated easily within minutes using these algorithms.

## **1.8 Structure of the Thesis**

Overall, this research is the study of chlorophyll *a* concentrations of phytoplankton using developed algorithms. Chapter 1 gives a small introduction to this study. This chapter also recounts the objectives and scope of the study. Chapter 2 describes the background of fresh water, phytoplankton, reflectance and chlorophyll *a* in details. Besides, in this chapter also provides graphs, images and tables on phytoplankton. Chapter 3 provides the publication of the theory of algorithms for measuring the concentration of chlorophyll *a* in the phytoplankton. This chapter also shows steps to develop the algorithms. In addition, techniques and measures are also given in this chapter. Chapter 4 provides research methods and tools that were used when the data were collected in this study. All measures carried out in this study were described in this chapter. Chapter 5 gives the data and also provides discussion of the study. This chapter also describes tests conducted on-site studies and other locations. Finally chapter 6 provides the overall conclusions of the study and provide recommendations for future research.

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 Introduction

Human is influenced strongly with aquatic ecosystem, especially phytoplankton. Everyone knows, phytoplankton gives more advantages in today's technology such as the measurement of phytoplankton biomass is important in aquatic ecology studies. It is frequently estimated from chlorophyll *a* concentration determination. The physical, chemical and biological factors (Platt & Denham, 1980; Smayda, 1980; Carrick et al, 1993; Cloern, 1996; Lucas et al, 1999a,b) are factors that influenced phytoplankton abundance in fresh water. Among the commonly discussed factors that influenced biomass loss are nutrient availability, control algal growth, light and temperature. The impacts of one factor are dependent on other the factors, for example, the nutrient loading effect on phytoplankton abundance in ecosystems is dependent on other factors, including effect biomass gains and losses, light availability (Hitchcock & Smayda, 1977; Cole & Cloern, 1984; Bledsoe & Philips, 2000), and sedimentation (Richardson & Jorgensen, 2013). The process of eutrophication can be accelerated (Smayda, 2013) in the area to significant human development. The frequency and intensity of algal bloom increasing can due to significant changes in the function and structure of the ecosystem effect, for example the water colour change of fresh water or ocean change from green to brown or from blue to red. Hence, in this chapter, definition and background of phytoplankton, reflectance and concentration of chlorophyll are given. Besides, the best method of chlorophyll *a* concentration measurement of phytoplankton is also discussed in this chapter.

## **2.2 Definition and Background of Phytoplankton**

Fresh water is a fundamental resource for human life, and the services provided by surface fresh water ecosystems underpin global water security, food security and economic productivity (Hanjra & Qureshi, 2010). Fresh water is defined as water having a low salt concentration which is usually less than 1%. Fresh water is vitally important to flora and fauna, including phytoplankton as a life resource. A good water quality of this system is required for many ecosystem services. Biological characteristics of fresh water, the development of flora and fauna in surface is governed by various species of animals and plants as well as physical performance of individual organism. The primary production of organic matter such as phytoplankton and macrophytes is extremely intensive in lakes and also reservoir but usually limited in rivers. Lakes have features low in average current velocity on surface value of 0.001 to 0.01 ms<sup>-1</sup> (Meybeck & Helmer, 1996).

Phytoplankton is also known as algae that heterogeneous assemblage of organisms. The first use of the term 'plankton' was by Viktor Hensen (Reynold 2006; Hutchinson, 1967; Schwartz 1968), German biologist in the latter half of the nineteenth century. He began a series of expeditions to gauge the abundance, composition and distribution of microscopic organism in the open ocean. However, the existence of such organism had been demonstrated by another investigator, Johannes Müller some years earlier. Table 2.1 shows the type of plankton based on different habitats.